SH
.U7F
no.698

NOAA Technical Report NMFS SSRF-698

Summer Benthic Fish Fauna of Sandy Hook Bay, New Jersey

STUART J. WILK and MYRON J. SILVERMAN

APR 14 1976
FISH & WILDLIFE SERVICE

SEATTLE, WA
January 1976

noaa NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION / National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries Series

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D83, Technical Information Division, Environmental Science Information Center, NOAA, Washington, D.C. 20235. Recent SSRF's are:

619. Macrozooplankton and small nekton in the coastal waters off Vancouver Island (Canada) and Washington, spring and fall of 1963. By Donald S. Day, January 1971, iii + 94 p., 19 figs., 13 tables.

620. The Trade Wind Zone Oceanography Pilot Study. Part IX: The sea-level wind field and wind stress values, July 1963 to June 1965. By Gunter R. Seckel. June 1970, iii + 66 p., 5 figs.

621. Predation by sculpins on fall chinook salmon, *Oncorhynchus tshawytscha*, fry of hatchery origin. By Benjamin G. Patten. February 1971, iii + 14 p., 6 figs., 9 tables.

622. Number and lengths, by season, of fishes caught with an otter trawl near Woods Hole, Massachusetts, September 1961 to December 1962. By F. E. Lux and F. E. Nichy. February 1971, iii + 15 p., 3 figs., 19 tables.

623. Apparent abundance, distribution, and migrations of albacore, *Thunnus alalunga*, on the North Pacific longline grounds. By Brian J. Rothschild and Marian Y. Y. Yong. September 1970, v + 37 p., 19 figs., 5 tables.

624. Influence of mechanical processing on the quality and yield of bay scallop meats. By N. B. Webb and F. B. Thomas. April 1971, iii + 11 p., 9 figs., 3 tables.

625. Distribution of salmon and related oceanograpic features in the North Pacific Ocean, spring 1968. By Robert R. French, Richard G. Bakkala, Masanao Osako, and Jun Ito. March 1971, iii + 22 p., 19 figs., 3 tables.

626. Commercial fishery and biology of the freshwater shrimp, *Macrobrachium*, in the Lower St. Paul River, Liberia, 1952-53. By George C. Miller. February 1971, iii + 13 p., 8 figs., 7 tables.

627. Calico scallops of the Southeastern United States, 1959-69. By Robert Cummins, Jr. June 1971, iii + 22 p., 23 figs., 3 tables.

628. Fur Seal Investigations, 1969. By NMFS, Marine Mammal Biological Laboratory. August 1971, 82 p., 20 figs., 44 tables, 23 appendix A tables, 10 appendix B tables.

629. Analysis of the operations of seven Hawaiian skipjack tuna fishing vessels, June-August 1967. By Richard N. Uchida and Ray F. Sumida. March 1971, v + 25 p., 14 figs., 21 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

630. Blue crab meat. I. Preservation by freezing. July 1971, iii + 13 p., 5 figs., 2 tables. II. Effect of chemical treatments on acceptability. By Jurgen H. Strasser, Jean S. Lennon, and Frederick J. King. July 1971, iii + 12 p., 1 fig., 9 tables.

631. Occurrence of thiaminase in some common aquatic animals of the United States and Canada. By R. A. Greig and R. H. Gnaedinger. July 1971, iii + 7 p., 2 tables.

632. An annotated bibliography of attempts to rear the larvae of marine fishes in the laboratory. By Robert C. May. August 1971, iii + 24 p., 1 appendix I table, 1 appendix II table. For sale by the Superintendent of Documents, U.S. Goverment Printing Office, Washington, D.C. 20402.

633. Blueing of processed crab meat. II Identification of some factors involved in the blue discoloration of canned crab meat *Callinectes sapidus*. By Melvin E. Waters. May 1971, iii + 7 p., 1 fig., 3 tables.

634. Age composition, weight, length, and sex of herring, *Clupea pallasii*, used for reduction in Alaska, 1929-66. By Gerald M. Reid. July 1971, iii + 25 p., 4 figs., 18 tables.

635. A bibliography of the blackfin tuna, *Thunnus atlanticus* (Lesson). By Grant L. Beardsley and David C. Simmons. August 1971, 10 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

636. Oil pollution on Wake Island from the tanker *R. C. Stoner*. By Reginald M. Gooding. May 1971, iii + 12 p., 8 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

637. Occurrence of larval, juvenile, and mature crabs in the vicinity of Beaufort Inlet, North Carolina. By Donnie L. Dudley and Mayo H. Judy. August 1971, iii + 10 p., 1 fig., 5 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

638. Length-weight relations of haddock from commercial landings in New England, 1931-55. By Bradford E. Brown and Richard C. Hennemuth. August 1971, v + 13 p., 16 figs., 6 tables, 10 appendix A tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

639. A hydrographic survey of the Galveston Bay system, Texas 1963-66. By E. J. Pullen, W. L. Trent, and G. B. Adams. October 1971, v + 13 p., 15 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

640. Annotated bibliography on the fishing industry and biology of the blue crab, *Callinectes sapidus*. By Marlin E. Tagatz and Ann Bowman Hall. August 1971, 94 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

641. Use of threadfin shad, *Dorosoma petenense*, as live bait during experimental pole-and-line fishing for skipjack tuna, *Katsuwonus pelamis*, in Hawaii. By Robert T. B. Iversen. August 1971, iii + 10 p., 3 figs., 7 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

642. Atlantic menhaden *Brevoortia tyrannus* resource and fishery—analysis of decline. By Kenneth A. Henry. August 1971, v + 32 p., 40 figs., 5 appendix figs., 3 tables, 2 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

643. Surface winds of the southeastern tropical Atlantic Ocean. By John M. Steigner and Merton C. Ingham. October 1971, iii + 20 p., 17 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

644. Inhibition of flesh browning and skin color fading in frozen fillets of yelloweye snapper (*Lutzanus vivanus*). By Harold C. Thompson, Jr., and Mary H. Thompson. February 1972, iii + 6 p., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

645. Traveling screen for removal of debris from rivers. By Daniel W. Bates, Ernest W. Murphey, and Martin G. Beam. October 1971, iii + 6 p., 6 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington D.C. 20402.

646. Dissolved nitrogen concentrations in the Columbia and Snake Rivers in 1970 and their effect on chinook salmon and steelhead trout. By Wesley J. Ebel. August 1971, iii + 7 p., 2 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington D.C. 20402.

647. Revised annotated list of parasites from sea mammals caught off the west coast of North America. By L. Margolis and M. D. Dailey. March 1972, iii + 23 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

Continued on inside back cover.

NOAA Technical Report NMFS SSRF-698

Summer Benthic Fish Fauna of Sandy Hook Bay, New Jersey

STUART J. WILK and MYRON J. SILVERMAN

SEATTLE, WA
January 1976

UNITED STATES
DEPARTMENT OF COMMERCE
Rogers C. B. Morton, Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Robert M. White, Administrator

National Marine
Fisheries Service
Robert W. Schoning, Director

For sale by the Superintendent of Documents, U.S. Government
Printing Office, Washington, D.C. 20402

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Page

Introduction 1
Description of study area 1
Materials and methods 1
Results 1
Summary 2
Literature cited 2

Figures

1. Outline of Sandy Hook Bay, N.J., divided into 18 blocks where finfish were sampled during an otter trawl survey, July to October 1970 2
2. The average catch (no.) of all fish per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 4
3. The average weight (kg) of all fish caught per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 4
4. The average catch (no.) of bay anchovy, *Anchoa mitchilli*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 4
5. The average catch (no.) of red hake, *Urophycis chuss*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 4
6. The average catch (no.) of spotted hake, *Urophycis regius*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 5
7. The average catch (no.) of scup, *Stenotomus chrysops*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 5
8. The average catch (no.) of weakfish, *Cynoscion regalis*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 5
9. The average catch (no.) of butterfish, *Peprilus triacanthus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 5
10. The average catch (no.) of northern searobin, *Prionotus carolinus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 6
11. The average catch (no.) of striped searobin, *Prionotus evolans*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 6
12. The average catch (no.) of windowpane, *Scophthalmus aquosus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 6
13. The average catch (no.) of winter flounder, *Pseudopleuronectes americanus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970 6
14. Monthly length-frequency distributions of red hake, *Urophycis chuss*, collected in Sandy Hook Bay, N.J., July to October 1970 7
15. Monthly length-frequency distributions of spotted hake, *Urophycis regius*, collected in Sandy Hook Bay, N.J., July to October 1970 7
16. Monthly length-frequency distributions of bluefish, *Pomatomus saltatrix*, collected in Sandy Hook Bay, N.J., July to October 1970 8
17. Monthly length-frequency distributions of scup, *Stenotomus chrysops*, collected in Sandy Hook Bay, N.J., July to October 1970 8
18. Monthly length-frequency distributions of weakfish, *Cynoscion regalis*, collected in Sandy Hook Bay, N.J., July to October 1970 9
19. Monthly length-frequency distributions of butterfish, *Peprilus triacanthus*, collected in Sandy Hook Bay, N.J., July to October 1970 9
20. Monthly length-frequency distributions of windowpane, *Scophthalmus aquosus*, collected in Sandy Hook Bay, N.J., July to October 1970 10
21. Monthly length-frequency distributions of winter flounder, *Pseudopleuronectes americanus*, collected in Sandy Hook Bay, N.J., July to October 1970 10

Table

1. Bottom temperature (T in °C), salinity (S in ppt), and depth (m) observed at otter trawl stations in Sandy Hook Bay, N.J., July to October 1970 3

Appendix Tables

1. List of species captured in order of total number caught by otter trawl in Sandy Hook Bay, N.J., July to October 1970 11
2. List of fishes captured by otter trawl in Sandy Hook Bay, N.J., July to October 1970, including summary data on time and location of capture, specimen size, and hydrographic characteristics at times of capture . 13

Summer Benthic Fish Fauna of Sandy Hook Bay, New Jersey

STUART J. WILK and MYRON J. SILVERMAN[1]

ABSTRACT

Thirty-eight species and 25 families of fishes were captured during an otter trawl survey in Sandy Hook Bay, N.J., during July to October 1970. Distribution, abundance, length and age composition, and environmental preferences were analyzed for the more numerous species. Winter flounder, *Pseudopleuronectes americanus*; striped searobin, *Prionotus evolans*; windowpane, *Scophthalmus aquosus*; and northern searobin, *Prionotus carolinus*, accounted for 68.3% by number and 66.4% by weight of the total survey catch.

INTRODUCTION

Historically, Sandy Hook Bay has supported commercial fisheries since the early days of settlement in the United States (True 1887). Within the past half century, water quality has deteriorated largely from sewage (Federal Water Pollution Control Administration 1967) and industrial waste contamination, and much of the bay bottom is now unfit for shellfish harvest. Only a few pound nets and purse seines are currently operated in the bay, but year-round and seasonally intensive small boat and shore recreational fisheries have developed. The following report of a systematic trawl survey of finfish is the first of several to be undertaken in Sandy Hook Bay, and provides baseline observations for future comparison of biological and environmental changes in the area.

Otter trawl hauls were made from July to October 1970 to determine summer species composition, relative abundance, and distribution of benthic fishes in Sandy Hook Bay. All catch data are summarized, together with length-frequency data and estimates of age composition and summer growth rates based on published age-length information.

DESCRIPTION OF STUDY AREA

Sandy Hook Bay is bounded on the north by Lower Bay, on the south and west by the New Jersey mainland, from which it receives drainage of the Navesink and Shrewsbury rivers, and on the east is separated from the Atlantic Ocean by Sandy Hook peninsula. To establish uniform sampling areas, the bay was divided into 18 blocks (Fig. 1). Except where interrupted by land, each block measured 1' lat. × 1' long., i.e., 1.8 km × 1.4 km (1.0 × 0.75 nautical mile).

MATERIALS AND METHODS

All 18 blocks were sampled, as close to biweekly as possible, during 2-day cruises when weather conditions and vessel scheduling permitted.

Our vessel was the Middle Atlantic Coastal Fisheries Center's *Martha-E II*, a 10.4-m (34-foot) diesel-powered research vessel equipped with hydraulic trawl winch, trawl boom, electronic navigation equipment, and electronic depth recorders.

At each sampling station, surface and bottom temperature and salinity were measured with a portable thermometer-salinometer probe. Station depth, bottom contours, and fish concentrations were observed with white-line fathometers aboard the vessel. The otter trawl was towed for 10 min at approximately 5.6 km/h (3 knots) in each sample block. The two-seam trawl used had the following dimensions: 9.1-m (30-foot) footrope and 7.6-m (25-foot) headrope; 64-mm (2.25-inch) stretched mesh 18-thread knotted nylon in the body of the trawl and 51-mm (2-inch) stretched mesh 18-thread knotless nylon in the cod end. During each trawl tow, frequent navigational checks were made to remain within a sample block. Although towing time remained constant, both direction and distance of the tow were governed by tide, current, wind, and starting location.

At the conclusion of each tow, the trawl was retrieved and emptied on the afterdeck of the vessel, where the fishes were sorted and identified. All specimens of each species were weighed and a random sample of 20-25 specimens measured to the nearest centimeter (total length). Usually, all specimens of each species were counted; but, when a species was very numerous, an estimate of the total was made by counting and weighing the fish in a subsample.

Physical and biological data collected were recorded on punch cards and subsequently sorted, mapped, and analyzed by computer.

Common and scientific names used follow those listed by the American Fisheries Society (Bailey et al. 1970).

RESULTS

During the 3.5-mo study, 112 station blocks were occupied on 15 sample dates. Data were grouped into eight sample periods of seven 2-day cruises and one 1-day cruise.

The results of our collections are presented as follows:

1. List of bottom temperature, salinity, and depth observed at each otter trawl station, including summary data on station blocks and sample periods (Table 1).
2. List of species caught in descending order of abundance, including summaries of each species and sample period (Appendix Table 1).
3. Phylogenetic list summarizing all parameters for each species (Appendix Table 2).
4. Maps of the average number and weight of all fish caught per 10-min tow (Figs. 2, 3).
5. Maps of the distribution of average catch (no.) per 10-min tow for 10 of the most numerous species (Figs. 4-13).

[1] Middle Atlantic Coastal Fisheries Center Sandy Hook Laboratory, National Marine Fisheries Service, NOAA, Highlands, NJ 07732.



Figure 1.—Outline of Sandy Hook Bay, N.J., divided into 18 blocks where finfish were sampled during an otter trawl survey, July to October 1970.

6. Length-frequency distributions representing the size distribution and changes in the size composition during the survey of eight species (Figs. 14-21).

SUMMARY

Four species—winter flounder, *Pseudopleuronectes americanus*; striped searobin, *Prinonotus evolans*; windowpane, *Scophthalmus aquosus*; and northern searobin, *Prionotus carolinus*—accounted for 68.3% by number and 66.4% by weight of the total catch during the survey. Although 10 of the most abundant species accounted for 95.0% by number and 84.6% by weight of the total catch, only seven species occurred in more than 25% of the collections (Appendix Table 1).

Catches in the northern half of Sandy Hook Bay (station blocks 1-9) contained a total of 35 species; those in the southern half (station blocks 10-18) 22 species (Appendix Table 2). The total catch both by weight and number in the northern half exceeded that of the southern half (Figs. 2, 3). The greater abundance and diversity of species in the northern half of the survey area were believed to be related to the deeper and slightly colder water found there (Table 1) and proximity to the ocean (Fig. 1).

The bay appears to provide a summer residence for several important recreational and commercial species, primarily red hake, *Urophycis chuss*; bluefish, *Pomatomus salatatrix*; scup, *Stenotomus chrysops*; weakfish, *Cynoscion regalis*; butterfish, *Peprilus triacanthus*; summer flounder, *Paralichthys dentatus*; and winter flounder, *Pseudopleuronectes americanus* (Appendix Table 2).

LITERATURE CITED

BAILEY, R. M., J. E. FITCH, E. S. HERALD, E. A. LACHNER, C. C. LINDSAY, C. R. ROBINS, and W. B. SCOTT.
1970. A list of common and scientific names of fishes from the United States and Canada. 3d ed. Am. Fish. Soc., Spec. Publ. 6, 149 p.

BARANS, C. A.
1972. Spotted hake, *Urophycis regius*, of the York River and lower Chesapeake Bay. Chesapeake Sci. 13:59-62.

BIGELOW, H. B., and W. C. SCHROEDER.
1953. Fishes of the Gulf of Maine. U.S. Fish Wildl. Serv., Fish. Bull. 53, 577 p.

FEDERAL WATER POLLUTION CONTROL ADMINISTRATION.
1967. Conference on pollution of Raritan Bay and adjacent interstate waters. Proc. 3d Sess., Vol. 1, N.Y., 448 p.

FINKELSTEIN, S. L.
1969. Age and growth of scup in the waters of eastern Long Island. N.Y. Fish Game J. 16:84-110.

LUX, F. E.
1973. Age and growth of the winter flounder, *Psudopleuronectes americanus*, on Georges Bank. Fish. Bull., U.S. 71:505-512.

McEACHRAN, J. D., and J. DAVIS.
1970. Age and growth of the striped searobin. Trans. Am. Fish. Soc. 99:343-352.

MOORE, E.
1947. Studies on the marine resources of southern New England. VI. The sand flounder, *Lophopsetta aquosa* (Mitchill); a general study of the species with special emphasis on age determination by means of scales and otoliths. Bull. Bingham Oceanogr. Collect. Yale Univ. 11:1-79.

NICHOLS, J. T., and C. M. BREDER, JR.
1927. The marine fishes of New York and southern New England. Zoologica (N.Y.) 9, 192 p.

Table 1.--Bottom temperature (T in °C), salinity (S in ppt), and depth (m) observed at otter trawl stations in Sandy Hook Bay, N. J., July to October 1970.

Station block		Sample period								Mean temp. and sal.	Mean depth
		July			August		September		October		
		2	16-17	30-31	13-14	27-28	10-11	24-25	15-16		
		1	2	3	4	5	6	7	8		
1	T	19.0	-	21.9	23.0	25.4	-	21.9	18.9	21.7	7.7
	S	26.2	-	26.1	27.4	26.8	-	28.8	26.3	26.9	
2	T	18.9	-	21.6	23.2	-	20.7	21.7	18.8	20.8	7.5
	S	26.1	-	26.3	27.7	-	27.8	27.8	26.8	27.1	
3	T	18.8	-	21.4	24.2	-	-	21.5	-	21.5	10.6
	S	26.8	-	26.8	27.6	-	-	29.4	-	27.7	
4	T	19.8	21.0	22.9	24.4	24.1	20.7	22.4	-	22.2	7.7
	S	26.0	26.4	26.2	27.3	26.7	28.1	28.2	-	27.0	
5	T	-	22.5	21.8	24.2	23.4	21.5	21.8	19.0	22.0	5.4
	S	-	25.3	25.9	26.8	27.0	27.0	27.7	27.1	26.7	
6	T	19.1	19.8	21.4	23.5	23.1	20.6	22.4	18.8	21.1	7.3
	S	26.5	27.1	26.7	27.5	27.5	28.1	28.3	26.2	27.2	
7	T	18.4	-	20.8	23.5	-	20.5	22.2	18.7	20.7	10.3
	S	27.3	-	27.0	28.0	-	28.5	28.8	26.6	27.7	
8	T	-	20.4	21.8	23.9	23.1	-	22.3	-	22.3	6.2
	S	-	26.6	26.6	27.6	27.6	-	29.5	-	27.6	
9	T	21.2	-	23.3	24.3	23.4	20.6	22.5	-	22.6	5.5
	S	25.0	-	26.3	27.6	26.8	28.3	28.6	-	27.1	
10	T	-	22.0	22.2	24.7	23.5	21.4	23.9	19.2	22.4	3.6
	S	-	25.3	26.0	26.7	26.5	26.7	26.4	25.6	26.2	
11	T	-	21.9	23.2	24.9	24.4	21.5	24.2	19.4	22.8	2.5
	S	-	25.6	25.8	26.8	25.8	27.0	26.6	25.7	26.2	
12	T	-	20.4	21.4	23.7	22.8	20.4	22.5	18.9	21.4	5.3
	S	-	26.6	27.0	27.4	27.2	28.5	28.0	26.1	27.3	
13	T	-	22.2	21.4	24.3	22.7	-	22.2	18.0	21.8	5.9
	S	-	26.0	27.0	27.1	27.5	-	28.2	26.0	27.0	
14	T	-	22.3	21.4	23.8	23.1	22.1	22.1	-	22.5	5.1
	S	-	26.0	26.9	27.5	27.3	27.5	27.7	-	27.2	
15	T	-	21.6	21.9	24.5	24.3	21.3	24.6	19.2	22.5	3.5
	S	-	25.8	26.1	26.8	26.0	27.6	26.1	25.9	26.3	
16	T	-	21.9	21.8	23.5	23.5	-	22.2	18.1	21.8	4.6
	S	-	25.8	26.7	27.4	26.9	-	27.2	26.5	26.8	
17	T	-	22.1	21.5	23.5	23.5	20.9	22.1	-	22.3	5.0
	S	-	26.2	26.9	27.5	26.7	28.0	27.7	-	27.2	
18	T	-	21.4	21.4	24.1	24.1	-	21.9	-	22.6	5.4
	S	-	26.3	27.0	27.4	26.0	-	27.7	-	26.9	
Mean	T	19.3	21.5	21.8	24.0	23.6	21.0	22.5	18.8	21.9	5.9
	S	26.3	26.1	26.5	27.3	26.8	27.8	27.9	26.3	27.0	

SMITH, R. W.
1969. An analysis of the summer flounder, *Paralichthys dentatus* (Linnaeus), population in the Delaware Bay. M.S. Thesis, Univ. Delaware, Newark, 72 p.

TRUE, F. W.
1887. The pound-net fisheries of the Atlantic States. *In* G. B. Goode (and a staff of Associates), The fisheries and fishery industries of the United States. Sect. V., p. 595-609. Gov. Print. Off., Wash. [D.C.]

WELSH, W. W., and C. M. BREDER, JR.
1924. Contributions to life histories of Sciaenidae of the eastern United States coast. Bull. U.S. Bur. Fish. 39:141-201.

WICKLUND, R. I., S. J. WILK, and L. OGREN.
1968. Observations on wintering locations of the northern pipefish and spotted seahorse. Underwater Nat. 5(2):26-28.

WONG, R. S. P.
1968. Age and growth of the northern searobin, *Prionotus carolinus* (Linnaeus). M.S. Thesis, Va. Inst. Mar. Sci., Gloucester Point, 48 p.



Figure 2.—The average catch (no.) of all fish per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 3.—The average weight (kg) of all fish caught per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 4.—The average catch (no.) of bay anchovy, *Anchoa mitchilli*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 5.—The average catch (no.) of red hake, *Urophycis chuss*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 6.—The average catch (no.) of spotted hake, *Urophycis regius*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 7.—The average catch (no.) of scup, *Stenotomus chrysops*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 8.—The average catch (no.) of weakfish, *Cynoscion regalis*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.

Figure 9.—The average catch (no.) of butterfish, *Peprilus triacanthus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 10.—The average catch (no.) of northern searobin, *Prionotus carolinus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 11.—The average catch (no.) of striped searobin, *Prionotus evolans*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 12.—The average catch (no.) of windowpane, *Scophthalmus aquosus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 13.—The average catch (no.) of winter flounder, *Pseudopleuronectes americanus*, per 10-min tow in Sandy Hook Bay, N.J., July to October 1970.



Figure 14.—Monthly length-frequency distributions of red hake, *Urophycis chuss*, collected in Sandy Hook Bay, N.J., July to October 1970.



Figure 15.—Monthly length-frequency distributions of spotted hake, *Urophycis regius*, collected in Sandy Hook Bay, N.J., July to October 1970.



Figure 16.—Monthly length-frequency distributions of bluefish, *Pomatomus saltatrix*, collected in Sandy Hook Bay, N.J., July to October 1970.



Figure 17.—Monthly length-frequency distributions of scup, *Stenotomus chrysops*, collected in Sandy Hook Bay, N.J., July to October 1970



Figure 18.—Monthly length-frequency distributions of weakfish, *Cynoscion regalis*, collected in Sandy Hook Bay, N.J., July to October 1970.



Figure 19.—Monthly length-frequency distributions of butterfish, *Peprilus triacanthus*, collected in Sandy Hook Bay, N.J., July to October 1970.



Figure 20.—Monthly length-frequency distributions of windowpane, *Scophthalmus aquosus*, collected in Sandy Hook Bay, N.J., July to October 1970.



Figure 21.—Monthly length-frequency distributions of winter flounder, *Pseudopleuronectes americanus*, collected in Sandy Hook Bay, N.J., July to October 1970.

Appendix Table 1.--List of species captured in order of total number caught by otter trawl in Sandy Hook Bay, N. J., July to October 1970.

Species	Sample period: July 2	July 16-17	July 30-31	August 13-14	August 27-28	September 10-11	September 24-25	October 15-16	Total	Percent of total catch	Percent occurrence	Average catch per 10-min tow
	1	2	3	4	5	6	7	8				
Winter flounder	29[1]	158	464	334	40	374	176	142	1,717	21.7	61.6	15.3
Pseudopleuronectes americanus	6.35[2]	28.87	84.99	40.72	3.18	56.65	31.33	19.00	271.09	24.8		2.42
Striped searobin	26	20	331	157	494	195	396	26	1,645	20.8	44.6	14.7
Prionotus evolans	2.72	3.29	55.63	21.83	81.56	39.50	74.25	4.65	283.43	26.0		2.53
Windowpane	78	32	161	96	4	119	326	234	1,050	13.3	45.5	9.4
Scophthalmus aquosus	13.15	4.82	22.00	6.66	0.12	14.35	32.49	20.64	114.23	10.5		1.02
Northern searobin	31	45	537	293	5	69	19	--	999	12.6	25.9	8.9
Prionotus carolinus	3.29	2.84	29.52	13.44	0.57	5.56	1.31	--	56.53	5.2		0.50
Weakfish	--	3	6	18	1	166	286	95	575	7.3	24.1	5.1
Cynoscion regalis	--	1.02	1.25	4.88	0.23	13.19	30.50	13.44	64.51	5.9		0.58
Bay anchovy	72	265	34	43	6	--	4	--	424	5.4	24.1	3.8
Anchoa mitchilli	0.68	1.59	0.19	0.28	0.02	--	0.02	--	2.78	0.3		0.02
Scup	17	8	9	64	63	27	133	100	421	5.3	37.5	3.8
Stenotomus chrysops	1.36	0.63	0.37	3.91	4.10	2.54	14.06	13.72	40.69	3.7		0.36
Butterfish	45	38	12	184	11	2	56	16	364	4.6	47.3	3.3
Peprilus triacanthus	4.58	1.59	0.52	4.63	0.15	0.06	1.16	0.69	13.38	1.2		0.12
Red hake	53	13	53	9	40	58	26	6	258	3.3	14.3	2.3
Urophycis chuss	3.29	0.60	3.69	0.79	4.82	8.16	4.88	1.81	28.04	2.6		0.25
Summer flounder	4	2	16	20	2	12	12	1	69	0.9	37.5	0.6
Paralichthys dentatus	4.65	1.13	9.41	13.50	2.23	7.14	9.64	0.91	48.61	4.5		0.43
Bluefish	2	9	5	--	--	8	3	37	64	0.8	24.1	0.6
Pomatomus saltatrix	0.99	1.13	2.04	--	--	1.31	0.91	4.54	10.92	1.0		0.10
Spotted hake	1	5	5	5	--	8	15	16	55	0.7	13.4	0.5
Urophycis regius	0.06	0.17	0.17	0.35	--	0.91	2.04	2.72	6.42	0.6		0.6
Grubby	8	21	2	--	1	9	2	--	43	0.5	7.1	0.4
Myoxocephalus aenaeus	0.17	0.79	0.17	--	0.07	0.34	0.06	--	1.60	0.1		0.01
Atlantic moonfish	--	2	--	--	2	9	8	15	36	0.5	15.2	0.3
Vomer setapinnis	--	0.05	--	--	0.03	0.11	0.07	0.15	0.41	*[3]		±[4]
Alewife	18	8	1	--	--	--	4	2	33	0.4	15.2	0.3
Alosa pseudoharengus	1.17	0.28	0.06	--	--	--	0.11	0.03	1.65	0.2		0.01
Northern kingfish	--	--	--	3	1	11	9	9	33	0.4	6.3	0.3
Menticirrhus saxatilis	--	--	--	0.62	0.23	1.76	1.36	1.07	5.04	0.5		0.05
American eel	--	1	--	--	1	14	11	2	29	0.4	6.3	0.3
Anguilla rostrata	--	0.79	--	--	0.53	8.96	6.80	0.91	17.99	1.6		0.16
Atlantic menhaden	1	1	--	1	1	6	10	--	20	0.3	11.6	0.2
Brevoortia tyrannus	0.34	0.23	--	0.45	0.23	2.04	3.29	--	6.58	0.6		0.06
Northern puffer	--	--	1	6	--	6	2	--	15	0.2	8.9	0.1
Sphoeroides maculatus	--	--	0.23	0.10	--	0.71	0.68	--	1.72	0.2		0.02
Atlantic sturgeon	--	1	6	--	--	--	7	--	14	0.2	2.7	0.1
Acipenser oxyrhynchus	--	1.47	12.59	--	--	--	38.56	--	52.62	4.8		0.47
Silver perch	--	--	--	1	--	10	--	2	13	0.2	4.5	0.1
Bairdiella chrysura	--	--	--	0.23	--	1.25	--	0.14	1.62	0.1		0.01
Tautog	2	--	--	1	--	1	5	--	9	0.1	4.5	0.1
Tautoga onitis	1.81	--	--	0.23	--	0.06	2.15	--	4.25	0.4		0.04
Sea raven	--	--	5	2	--	--	--	--	7	0.1	1.8	0.1
Hemitripterus americanus	--	--	0.15	0.11	--	--	--	--	0.26	*		±
Shortnose sturgeon	--	1	3	--	--	2	--	--	6	0.1	2.7	0.1
Acipenser brevirostrum	--	5.44	19.28	--	--	15.88	--	--	40.60	3.7		0.36

Appendix Table 1.--Continued.

Species	Sample period: July 2	July 16-17	July 30-31	August 13-14	August 27-28	September 10-11	September 24-25	October 15-16	Total	Percent of total catch	Percent occurrence	Average catch per 10-min tow
	1	2	3	4	5	6	7	8				
Roughtail stingray	--	2	1	--	--	1	1	--	5	0.1	4.5	**[5]
Dasyatis centroura	--	4.99	2.27	--	--	2.04	3.63	--	12.93	1.2		0.10
American shad	--	--	1	--	--	--	1	--	2	*	1.8	**
Alosa sapidissima	--	--	0.23	--	--	--	0.23	--	0.46	*		±
Conger eel	--	--	--	--	--	1	--	1	2	*	1.8	**
Conger oceanicus	--	--	--	--	--	0.34	--	0.03	0.37	*		±
Smallmouth flounder	--	--	--	--	--	--	1	1	2	*	1.8	**
Etropus microstomus	--	--	--	--	--	--	0.03	0.01	0.04	*		±
Oyster toadfish	--	--	--	--	--	--	2	--	2	*	1.8	**
Opsanus tau	--	--	--	--	--	--	0.17	--	0.17	*		±
Planehead filefish	--	--	--	--	--	--	2	--	2	*	0.9	**
Monacanthus hispidus	--	--	--	--	--	--	0.06	--	0.06	*		±
Atlantic silverside	--	--	--	--	--	1	--	--	1	*	0.9	**
Menidia menidia	--	--	--	--	--	0.01	--	--	0.01	*		±
Blueback herring	1	--	--	--	--	--	--	--	1	*	0.9	**
Alosa aestivalis	0.03	--	--	--	--	--	--	--	0.03	*		±
Clearnose skate	--	--	1	--	--	--	--	--	1	*	0.9	**
Raja eglanteria	--	--	1.36	--	--	--	--	--	1.36	0.1		0.01
Lined seahorse	--	--	--	--	--	--	--	1	1	*	0.9	**
Hippocampus erectus	--	--	--	--	--	--	--	0.03	0.03	*		±
Lookdown	--	--	1	--	--	--	--	--	1	*	0.9	**
Selene vomer	--	--	0.01	--	--	--	--	--	0.01	*		±
Orange filefish	--	--	--	--	--	--	--	1	1	*	0.9	**
Aluterus schoepfi	--	--	--	--	--	--	--	0.57	0.57	0.1		0.01
Smooth dogfish	--	--	1	--	--	--	--	--	1	*	0.9	**
Mustelus canis	--	--	0.91	--	--	--	--	--	0.91	0.1		0.01
Striped burrfish	--	--	--	--	--	--	--	1	1	*	0.9	**
Chilomycterus schoepfi	--	--	--	--	--	--	--	0.06	0.06	*		±
Total	388	635	1,656	1,237	672	1,109	1,517	708	7,922			
	44.64	61.72	247.04	112.73	98.07	182.87	259.79	85.12	1,091.98			
Number of tows	7	13	18	18	15	12	18	11	112			
Average catch	55.4	48.8	92.0	68.7	44.8	92.4	84.3	64.4	70.7			
per 10-min tow	6.38	4.75	13.72	6.26	6.54	15.24	14.43	7.74	9.75			

[1]Number of fish.
[2]Weight of fish in kilograms.
[3]Less than 0.1% of total catch.
[4]Less than 0.01 kg per 10-min tow.
[5]Less than 0.1% fish per 10-min tow.

Appendix Table 2.--List of fishes captured by otter trawl in Sandy Hook Bay, N. J., July to October 1970, including summary data on time and location of capture, specimen size, and hydrographic characteristics at times of capture.

Species	Occurrence: Month(s)	Occurrence: Station block(s)	Total length: Range	Total length: Mean	Bottom temperature: Range	Bottom temperature: Mean	Bottom salinity: Range	Bottom salinity: Mean	Depth: Range	Depth: Mean	Remarks
			cm	cm	°C	°C	ppt	ppt	m	m	
Smooth dogfish *Mustelus canis*	July	4	--	62.0	--	22.9	--	26.2	--	7.0	
Clearnose skate *Raja eglanteria*	July	4	--	57.0	--	22.9	--	26.2	--	7.0	
Roughtail stingray *Dasyatis centroura*	July	4,13,17	80-100	90.0	21.0-22.2	21.6	26.0-26.9	26.4	5.5- 8.2	6.4	
	Sept.	7,10	--	100.0	21.4-22.2	21.8	26.7-28.8	27.8	3.0- 7.6	5.3	
Shortnose sturgeon *Acipenser brevoortia*	July	4	80-100	92.0	21.0-22.9	22.0	26.2-26.4	26.3	7.0- 8.2	7.6	Ninety percent of all sturgeon species were taken in block 4, the area of the survey most affected by currents, tides, and proximity to the ocean and other bay systems.
	Sept.	6	92-132	112.0	--	20.6	--	28.1	--	5.8	
Atlantic sturgeon *Acipenser oxyrhynchus*	July	4	70- 96	86.0	21.0-22.9	22.0	26.2-26.4	26.3	7.0- 8.2	7.6	
	Sept.	4	100-150	135.0	--	22.4	--	28.2	--	11.6	
American eel *Anguilla rostrata*	July	4	--	65.0	--	21.0	--	26.4	--	8.2	This species seemed to prefer the cooler and deeper waters found in the northern half of the bay.
	Aug.	11	--	67.0	--	24.4	--	25.8	--	3.0	
	Sept.	4,6,10	40- 55	48.3	20.6-22.4	21.3	26.7-28.3	27.8	3.0- 6.7	5.3	
	Oct.	10	41- 45	43.0	--	19.2	--	25.6	--	3.7	
Conger eel *Conger oceanicus*	Sept.	4	--	32.0	--	20.7	--	28.1	--	6.7	
	Oct.	6	--	17.0	--	18.8	--	26.2	--	9.1	
Blueback herring *Alosa aestivalis*	July	7	--	14.0	--	18.4	--	27.3	--	10.4	Herring species can only be considered incidental in our trawl catches because of their pelagic nature.
Alewife *Alosa pseudoharengus*	July	1,2,4,6,7	12- 28	14.6	18.4-22.9	19.8	26.1-27.3	26.6	7.0-10.4	8.6	
	Sept.	7	16- 18	17.0	--	22.2	--	28.8	--	7.6	
	Oct.	2	--	18.0	--	18.8	--	26.8	--	8.5	
American shad *Alosa sapidissima*	July	10	--	28.0	--	22.2	--	26.0	--	4.6	
	Sept.	4	--	34.0	--	22.4	--	28.2	--	11.6	
Atlantic menhaden *Brevoortia tyrannus*	July	9,13	29- 30	29.5	21.2-22.2	21.7	25.0-26.0	25.5	2.7- 5.5	4.1	
	Aug.	8,11	33- 39	36.0	23.9-24.4	24.2	25.8-27.6	26.7	3.0- 6.1	4.6	
	Sept.	1,4,6,7, 10,12, 14,15	30- 39	33.1	20.4-22.4	21.6	26.7-28.8	28.1	2.4-11.6	6.4	
Bay anchovy *Anchoa mitchilli*	July	1-3,5,6, 9-13,15, 17,18	6- 9	7.6	18.8-23.3	20.9	25.3-27.1	26.3	2.1-10.1	6.1	Most caught in southwest corner of the bay (Fig. 4).
	Aug.	1,2,6, 10-12, 14-16,18	7- 9	7.8	23.0-24.9	23.9	25.8-27.7	27.1	2.4- 6.7	5.0	Judging from stomach contents of bluefish and striped bass, the bay anchovy is a major food item of these higher predators.
	Sept.	1,7	6- 8	6.8	21.9-22.2	22.1	--	28.8	6.7- 7.6	7.2	
Oyster toadfish *Opsanus tau*	Sept.	16,18	14- 20	17.0	21.9-22.2	22.1	27.2-27.7	27.5	4.6- 5.2	4.9	

Appendix Table 2.--Continued.

Species	Occurrence: Month(s)	Occurrence: Station block(s)	Total length: Range	Total length: Mean	Bottom temperature: Range	Bottom temperature: Mean	Bottom salinity: Range	Bottom salinity: Mean	Depth: Range	Depth: Mean	Remarks
			cm	cm	°C	°C	ppt	ppt	m	m	
Red hake Urophycis chuss	July	1-4,6,7	11- 28	18.0	18.4-22.9	20.8	26.0-27.3	26.6	7.0-12.2	9.0	All taken in northern half of bay 96% in blocks 1, 4, and 7 (Fig. 5)
	Aug.	1,7	16- 27	23.4	23.5-25.4	24.5	26.8-28.0	27.4	10.1-12.2	11.2	
	Sept.	4,7	19- 32	25.2	20.5-22.4	21.5	28.1-28.8	28.4	6.7-11.6	9.2	
	Oct.	1	29- 36	31.0	--	18.8	--	26.2	--	9.1	According to age-length data (Bigelow and Schroeder 1953), the majority of our catch (Fig. 14) was made up of yearlings.
Spotted hake Urophycis regius	July	3,4,8	12- 20	15.5	19.8-22.9	21.1	26.0-26.8	26.4	5.5-12.2	8.0	All but one taken in northern half of survey area; 74.5% in blocks 4 and 7 (Fig. 6).
	Aug.	3,7,12	17- 18	17.2	23.5-24.2	23.8	27.4-28.0	27.7	5.5-12.2	8.9	
	Sept.	3,4,7	20- 26	23.0	20.5-22.4	21.5	28.1-29.4	28.6	6.7-13.4	10.1	
	Oct.	6,7	21- 27	25.1	18.7-18.8	18.8	26.2-26.6	26.4	9.1- 9.8	9.5	Catch was composed primarily of young-of-the-year specimens, as indicated by monthly length-frequencies (Fig. 15) and growth data given by Barans (1972).
Atlantic silverside Menidia menidia	Sept.	11	--	6.0	--	21.5	--	27.0	--	1.8	Judging from previous examination of beach seine catches, this shore species is one of the most numerou in the survey area. From its lack of appearance in our trawl catch, it is apparent gear selectivity and the inshore habit of this species limited catches.
Lined seahorse Hippocampus erectus	Oct.	10	--	14.0	--	19.4	--	25.7	--	2.4	Year-round resident in New York Bight (Wicklund, Wilk, and Ogren, 1968).
Bluefish Pomatomus saltatrix	July	2,3,5, 10-15	7- 39	23.7	18.8-23.2	21.4	25.3-26.8	26.1	2.1-12.2	5.8	From age determinations based on scales (Wilk, unpublished data), 76.5% of our catch (Fig. 16) were young-of-the-year specimens.
	Sept.	1,2,4,5, 9,10,12, 15,17	17- 24	17.6	20.4-22.4	21.2	26.7-28.8	27.9	3.0-11.6	6.1	
	Oct.	1,2,5,6, 12,13,16	19- 43	24.6	18.0-19.0	18.6	26.0-27.1	26.4	4.0- 9.1	7.2	
Lookdown Selene vomer	July	6	--	4.0	--	21.4	--	26.7	--	7.3	
Atlantic moonfish Vomer setapinnis	July	14,17	5- 6	5.5	22.1-22.3	22.2	26.0-26.2	26.1	4.3- 5.5	4.9	Both the lookdown and Atlantic moonfish are considered only summer visitors from southern waters (Bigelow and Schroeder 1953, and Nichols and Breder 192
	Aug.	8	--	6.0	--	23.1	--	27.6	--	5.8	
	Sept.	1,5-7,9, 10	4- 8	7.1	20.6-22.4	21.5	26.7-28.8	28.0	3.0- 7.6	5.8	
	Oct.	1,2,5,7, 10,13,16	5- 8	6.0	18.0-19.2	18.7	25.6-27.1	26.4	3.7- 9.8	7.2	
Scup Stenotomus chrysops	July	4,6,8,10	10- 20	13.4	19.8-22.9	21.3	25.3-26.7	26.3	2.7- 8.5	6.6	Specimens taken in all blocks except 14, 16, 17, and 18; larges average catch in block 7 (21.8 pe
	Aug.	1-6,8-11, 13	11- 18	14.9	22.7-25.4	23.9	25.8-27.7	27.2	2.4-10.1	5.9	
	Sept.	2,4-7, 9-12,15	13- 20	16.8	20.4-22.4	21.2	26.7-28.8	27.9	1.8-11.6	6.3	10-min tow) (Fig. 7).
	Oct.	2,6,7,10, 13	16- 21	18.9	18.0-19.2	18.7	25.6-26.8	26.2	3.7- 9.8	7.7	Comparing monthly length-frequencies (Fig. 17) to age-leng data given by Bigelow and Schroed (1953) and Finkelstein (1969), th majority of the scup caught were probably yearlings.
Silver perch Bairdiella chrysura	Aug.	10	--	21.0	--	24.3	--	27.6	--	6.7	
	Sept.	4,12	10- 20	18.6	20.4-20.7	20.6	28.1-28.5	28.3	5.5- 6.7	6.1	
	Oct.	6,7	12- 22	17.0	18.7-18.8	18.8	26.2-26.6	26.4	9.1- 9.8	9.5	

Appendix Table 2.--Continued.

Species	Occurrence: Month(s)	Occurrence: Station block(s)	Total length: Range	Total length: Mean	Bottom temperature: Range	Bottom temperature: Mean	Bottom salinity: Range	Bottom salinity: Mean	Depth: Range	Depth: Mean	Remarks
			cm	cm	°C	°C	ppt	ppt	m	m	
Weakfish	July	7,10,11	24- 39	27.7	20.8-22.2	21.6	25.6-27.0	26.2	2.1-10.7	5.8	Most (82.6%) were taken in northern
Cynoscion regalis	Aug.	3,7-11,16	11- 46	27.9	23.5-24.9	24.2	25.8-28.0	27.2	2.4-12.2	6.1	half of survey area. Blocks 6 and
	Sept.	4-10,12	11- 51	18.2	20.5-22.5	21.5	26.7-29.5	28.2	3.0-11.6	6.6	7 combined accounted for 55.8% of
	Oct.	2,6,7,13	15- 51	19.8	18.0-18.8	18.6	26.0-26.8	26.4	7.6- 9.8	8.8	all weakfish caught (Fig. 8).
											Comparing monthly length-frequencies (Fig. 18) to growth information given by Welsh and Breder (1924) and Bigelow and Schroeder (1953), the majority of the weakfish were probably 1- and 2-year-old fish; mostly the former.
Northern kingfish	Aug.	1,7,8,11	12- 30	24.8	23.0-24.4	23.7	25.8-28.0	27.2	3.0-12.2	6.9	All but three northern kingfish
Menticirrhus saxatilis	Sept.	4,6,7,9, 10	17- 34	23.3	20.5-22.5	21.5	26.7-28.8	28.2	3.0-11.6	7.1	were caught in northern half of bay.
	Oct.	1,6,10,16	11- 31	16.3	18.1-19.2	18.8	25.6-26.5	26.2	3.7- 9.1	7.0	
Tautog	July	4	23- 37	30.0	--	18.8	--	26.8	--	7.3	
Tautoga onitis	Aug.	10	--	19.0	--	24.7	--	26.7	--	3.7	
	Sept.	4,6	18- 26	19.2	20.7-22.4	21.8	28.1-28.3	28.2	5.8-11.6	8.0	
Butterfish	July	1-3,5-8, 10,11, 13-18	3- 24	14.1	18.4-22.5	21.0	25.3-27.3	26.3	2.7-10.7	5.7	Butterfish were taken in every
Peprilus triacanthus	Aug.	1,2,5-16, 18	4- 23	10.5	22.7-24.9	23.8	26.0-28.0	27.2	2.4-12.2	5.6	block except 4; highest average catches in blocks 1, 2, and 11 (Fig. 9).
	Sept.	1-3,6-8,12	8- 19	11.7	20.4-22.4	21.8	28.7-29.5	28.7	4.9-13.4	7.3	
	Oct.	1,6,7,16	11- 23	15.1	18.1-18.9	18.6	26.2-26.6	26.4	6.7- 9.8	8.5	Judging from monthly length-frequencies (Fig. 19) and growth data given by Bigelow and Schroeder (1953), the majority of butterfish taken were in their second and third summers, with the yearlings dominating the catch.
Northern searobin	July	2-4,6-9, 12,14	12- 27	19.0	18.4-23.3	21.2	25.0-27.3	26.5	2.7-12.2	7.6	Almost all (99.1%) were taken in
Prionotus carolinus	Aug.	3,4,6-9, 13,15	15- 20	17.5	23.1-24.5	24.0	26.8-28.0	27.4	3.7-12.2	6.9	northern half of survey area; 62.5% in blocks 3 and 4 (Fig. 10).
	Sept.	3,4,6,7	17- 21	19.4	20.5-22.4	21.3	28.1-29.4	28.5	5.8-13.4	9.4	According to growth data given by Wong (1968), our specimens were 1- to 7-year-old fish.
Striped searobin	July	2-4,6-10, 12-17	15- 30	20.0	18.4-23.3	21.3	25.0-27.3	26.5	2.7-12.2	6.8	Striped searobin were taken in all
Prionotus evolans	Aug.	1-9,11,12, 14,16	12- 22	17.7	22.8-25.4	23.8	25.8-28.0	27.3	3.0-12.2	6.6	except block 18; 90.3% in northern half of survey area (Fig. 11).
	Sept.	1-8,10-12	9- 36	21.1	20.4-24.2	21.6	26.6-29.5	28.1	1.8-13.4	6.7	Blocks 6-8 accounted for 64.0%
	Oct.	1,2,5-7, 13-15	10- 35	19.5	18.7-19.2	18.9	25.9-27.1	26.4	2.4- 9.8	7.4	of this species total catch.
											On the basis of age-length data given by McEachran and Davis (1970), it appears that our samples were composed of all ages from young-of-the-year to 6-year-old fish.
Sea raven	July	3	11- 12	11.5	--	21.9	--	26.1	--	7.6	
Hemitripterus americanus	Aug.	4	10- 15	12.5	--	24.4	--	27.3	--	6.1	
Grubby	July	4,7	9- 18	11.6	18.4-22.9	20.6	26.0-27.3	26.6	7.0-10.7	8.7	All specimens taken in northern
Myoxocephalus aenaeus	Aug.	1	--	10.0	--	25.4	--	26.8	--	10.1	half of survey area; 81.4% in
	Sept.	4	9- 18	11.5	20.7-22.4	21.6	28.1-28.2	28.2	6.7-11.6	9.2	block 4.

Appendix Table 2.--Continued.

Species	Occurrence: Month(s)	Occurrence: Station block(s)	Total length: Range	Total length: Mean	Bottom temperature: Range	Bottom temperature: Mean	Bottom salinity: Range	Bottom salinity: Mean	Depth: Range	Depth: Mean	Remarks
			cm	cm	°C	°C	ppt	ppt	m	m	
Smallmouth flounder	Sept.	5	--	12.0	--	21.8	--	27.7	--	4.6	
Etropus microstomus	Oct.	6	--	6.0	--	18.8	--	26.2	--	9.1	
Summer flounder	July	2-4,6-10, 12-16	32- 49	39.5	18.4-23.3	21.5	25.0-27.3	26.4	2.7-12.2	6.9	Judging from growth data given by Smith (1969), our samples were composed of fish 2-5 years old. This species is one of the most sought after by sportsmen in the area.
Paralichthys dentatus	Aug.	1-3,5-9, 11-13	35- 51	40.3	23.0-25.4	24.0	25.8-28.0	27.2	2.4-12.2	6.4	
	Sept.	2-7,9,12	29- 47	39.4	20.4-22.4	21.3	27.0-29.4	28.2	5.2-13.4	7.5	
	Oct.	5	--	40.0	--	19.0	--	27.1	--	5.8	
Windowpane	July	2-8,12, 13,15	7- 31	21.7	18.4-22.9	21.0	25.9-27.3	26.6	4.3-12.2	7.8	Most windowpane (97.8%) were taken in northern half of survey area; 85.6% in blocks 3, 4, and 7 (Fig. 12).
Scophthalmus aquosus	Aug.	1-9,13, 14,16	7- 28	14.6	22.7-25.4	23.8	26.8-28.0	27.4	4.6-12.2	6.5	
	Sept.	1-4,6,7, 14	9- 29	18.5	20.5-22.4	21.5	27.5-29.4	28.3	2.4- 9.8	7.0	Based on growth data given by Moore (1947) and monthly length-frequencies (Fig. 20), July catches were probably composed mainly of 4- and 5-year-old fish. In later months, 2- and 3-year olds predominated.
	Oct.	1,2,5-7, 12,13,15	12- 25	18.2	18.0-19.2	18.8	25.9-27.1	26.4	2.4- 9.8	7.0	Windowpane preferred the cooler and deeper waters found in the northern half of the survey area. All catches of more than 35 specimens, i.e., 77.7% of the total catch, were made in water which averaged nearly 4 m deeper and 1°C cooler than the survey average.
Winter flounder	July	2-9,11-18	10- 37	21.3	18.4-22.2	21.3	25.8-27.3	26.5	2.4-10.7	6.4	Winter flounder occurred in all blocks. Heaviest catches in northern half of bay; 78.2% in blocks 3, 4, 6, and 7 (Fig. 13).
Pseudopleuronectes americanus	Aug.	1-9,11-14	11- 34	20.8	23.0-24.9	23.8	26.8-28.0	27.4	2.4-12.2	6.3	
	Sept.	1-4,6-10, 12,14	10- 36	21.9	20.4-22.5	21.5	26.7-29.5	28.3	2.4-13.4	7.0	
	Oct.	1,5-7, 10-13, 15,16	11- 32	20.2	18.0-19.4	18.8	25.6-27.1	26.2	2.4- 9.8	6.0	Judging from data given by Bigelow and Schroeder (1953) and Lux (1973) and monthly length-frequencies (Fig. 21), the measured specimens were 1-4 years of age.
											Catches of over 100 individuals were all made in the 6.7- to 11.0-m depth range; the mean depth in which these larger catches were made is 2.8 m greater than the survey average.
Orange filefish	Oct.	6	--	32.0	--	18.8	--	26.2	--	9.1	Both species of filefish are rather common in general area during summer and fall (Nichols and Breder 1927).
Aluterus schoepfi											
Planehead filefish	Sept.	4	10- 11	10.5	--	22.4	--	28.2	--	11.6	
Monocanthus hispidus											
Northern puffer	July	4	--	21.0	--	22.9	--	26.2	--	7.0	All specimens taken in northern half of survey area.
Sphoeroides maculatus	Aug.	4-7	6- 8	7.0	23.5-24.4	23.9	26.8-28.0	27.4	3.7-12.2	7.2	
	Sept.	4,6,7,9	9- 27	18.5	20.6-22.4	21.3	28.1-28.8	28.3	5.8-11.6	7.3	
Striped burrfish	Oct.	6	--	10.0	--	18.8	--	26.2	--	9.1	This species, according to Bigelow and Schroeder (1953) and Nichols and Breder (1927), is an occasional summer visitor to New Jersey waters
Chilomycterus schoepfi											

★ U.S. GOVERNMENT PRINTING OFFICE

648. **Weight loss of pond-raised channel catfish (*Ictalurus punctatus*) during holding in processing plant vats.** By Donald C. Greenland and Robert L. Gill. December 1971, iii + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

649. Distribution of forage of skipjack tuna (*Euthynnus pelamis*) in the eastern tropical Pacific. By Maurice Blackburn and Michael Laurs. January 1972, iii + 16 p., 7 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

650. Effects of some antioxidants and EDTA on the development of rancidity in Spanish mackerel (*Scomberomorus maculatus*) during frozen storage. By Robert N. Farragut. February 1972, iv + 12 p., 6 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

651. The effect of premortem stress, holding temperatures, and freezing on the biochemistry and quality of skipjack tuna. By Ladell Crawford. April 1972, iii + 23 p., 3 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

653. The use of electricity in conjunction with a 12.5-meter (Headrope) Gulf-of-Mexico shrimp trawl in Lake Michigan. By James E. Ellis. March 1972, iv + 10 p., 11 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

654. An electric detector system for recovering internally tagged menhaden, genus *Brevoortia*. By R. O. Parker, Jr. February 1972, iii + 7 p., 3 figs., 1 appendix table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

655. Immobilization of finerling salmon and trout by decompression. By Doyle F. Sutherland. March 1972, iii + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

656. The calico scallop, *Argopecten gibbus*. By Donald M. Allen and T. J. Costello. May 1972, iii + 19 p., 9 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

657. Making fish protein concentrates by enzymatic hydrolysis. A status report on research and some processes and products studied by NMFS. By Malcolm B. Hale. November 1972, v + 32 p., 15 figs., 17 tables, 1 appendix table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

658. List of fishes of Alaska and adjacent waters with a guide to some of their literature. By Jay C. Quast and Elizabeth L. Hall. July 1972, iv + 47 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

659. The Southeast Fisheries Center bionumeric code. Part I: Fishes. By Harvey R. Bullis, Jr., Richard B. Roe, and Judith C. Gatlin. July 1972, xi + 95 p., 2 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

660. A freshwater fish electro-motivator (FFEM)-its characteristics and operation. By James E. Ellis and Charles C. Hoopes. November 1972, iii + 11 p., 9 figs.

661. A review of the literature on the development of skipjack tuna fisheries in the central and western Pacific Ocean. By Frank J. Hester and Tamio Otsu. January 1973, iii + 13 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

662. Seasonal distribtution of tunas and billfishes in the Atlantic. By John P. Wise and Charles W. Davis. January 1973, iv + 24 p., 13 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

663. Fish larvae collected from the northeastern Pacific Ocean and Puget Sound during April and May 1967. By Kenneth D. Waldron. December 1972, iii + 16 p., 2 figs., 1 table, 4 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington D.C. 20402.

664. Tagging and tag-recovery experiments with Atlantic menhaden, *Brevoortia tyrannus*. By Richard L. Kroger and Robert L. Dryfoos. December 1972, iv + 11 p., 4 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

665. Larval fish survey of Humbolt Bay, California. By Maxwell B. Eldrige and Charles F. Bryan. December 1972, iii + 8 p., 8 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

666. Distribution and relative abundance of fishes in Newport River, North Carolina. By William R. Turner and George N. Johnson. September 1973, iv + 23 p., 1 fig., 13 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

667. An analysis of the commercial lobster (*Homarus americanus*) fishery along the coast of Maine, August 1966 through December 1970. By James C. Thomas. June 1973, v + 57 p., 18 figs., 11 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

668. An annotated bibliography of the cunner, *Tautogolabrus adspersus* (Walbaum). By Fredric M. Serchuk and David W. Frame. May 1973, ii + 43 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

669. Subpoint prediction for direct readout meterological satellites. By L. E. Eber. August 1973, iii + 7 p., 2 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

670. Unharvested fishes in the U.S. commercial fishery of western Lake Erie in 1969. By Harry D. Van Meter. July 1973, iii + 11 p., 6 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

671. Coastal upwelling indices, west coast of North America, 1946-71. By Andrew Bakun. June 1973, iv + 103 p., 6 figs., 3 tables, 45 appendix figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

672. Seasonal occurrence of young Gulf menhaden and other fishes in a northwestern Florida estuary. By Marlin E. Tagatz and E. Peter H. Wilkins. August 1973, iii + 14 p., 1 fig., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

673. Abundance and distribution of inshore benthic fauna off southwestern Long Island, N.Y. By Frank W. Steimle, Jr. and Richard B. Stone. December 1973, iii + 50 p., 2 figs., 5 appendix tables.

674. Lake Erie bottom trawl explorations, 1962-66. By Edgar W. Bowman. January 1974, iv + 21 p., 9 figs., 1 table, 7 appendix tables.

675. Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972. Part 2. Review and Contributed Papers. Richard S. Shomura and Francis Williams (editors). July 1974, iv + 335 p., 38 papers. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

676. Price spreads and cost analyses for finfish and shellfish products at different marketing levels. By Erwin S. Penn. March 1974, vi + 74 p., 15 figs., 12 tables, 12 appendix figures, 41 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

677. Abundance of benthic macroinvertebrates in natural and altered estuarine areas. By Gill Gilmore and Lee Trent. April 1974, iii + 13 p., 11 figs., 3 tables, 2 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

678. Distribution, abundance, and growth of juvenile sockeye salmon, *Oncorhynchus nerka*, and associated species in the Naknek River system, 1961-64. By Robert J. Ellis. September 1974, v + 53 p., 27 figs., 26 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

679. Kinds and abundance of zooplankton collected by the USCG icebreaker *Glacier* in the eastern Chukchi Sea, September-October 1970. By Bruce L. Wing, August 1974, iv + 18 p., 14 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

680. Pelagic amphipod crustaceans from the southeastern Bering Sea, June 1971. By Gerald A. Sanger. July 1974, iii + 8 p., 3 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

681. Physiological response of the cunner, *Tautogolabrus adspersus*, to cadmium. October 1974, iv + 33 p., 6 papers, various authors. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

682. Heat exchange between ocean and atmosphere in the eastern North Pacific for 1961-71. By N. E. Clark, L. Eber, R. M. Laurs, J. A. Renner, and J. F. T. Saur. December 1974, iii + 108 p., 2 figs., 1 table, 5 plates.

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS STAFF
ROOM 450
1107 N.E. 45TH ST.
SEATTLE, WA 98105

OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-210
THIRD CLASS
BULK RATE

U.S. MAIL



AMERICAN REVOLUTION BICENTENNIAL 1776-1976

76 AMERICA'S FIRST INDUSTRY